arXiv:quant-ph/0612210v3 22 Jun 2007

Collective multipole-like signatures of entanglement in symmetric N-qubit systems

A. R. Usha Devi,[1, *] R. Prabhu,[1] and A. K. Rajagopal[2]

[1]*Department of Physics, Bangalore University, Bangalore-560 056, India*
[2]*Department of Computer Science and Center for Quantum Studies, George Mason University, Fairfax, VA 22030, USA and Inspire Institute Inc., McLean, VA 22101, USA.*

(Dated: February 1, 2008)

A cogent theory of *collective multipole-like quantum correlations* in symmetric multiqubit states is presented by employing SO(3) irreducible spherical tensor representation. An *arbitrary bipartite division* of this system leads to a family of inequalities to detect entanglement involving averages of these tensors expressed in terms of the total system angular momentum operator. Implications of this theory to the quantum nature of multipole-like correlations of all orders in the Dicke states are deduced. A selected set of examples illustrate these collective tests. Such tests detect entanglement in macroscopic atomic ensembles, where individual atoms are not accessible.

PACS numbers: 03.67.-a, 03.67.Mn, 03.65.-w, 03.65.Ca

Correlated macroscopic atomic ensembles [1] offer promising possibilities in low-noise spectroscopy [2], high precision interferometry [3], and in the implementation of quantum information protocols [4]. Experimental characterization of entanglement - which is the key ingredient in these applications - has attracted considerable attention. Main difficulty in analyzing such composite systems with large number, N, of particles is the corresponding exponential size of the Hilbert space. So, major effort is focused on exploring inseparability status of special classes [5, 6] of quantum states, confined to smaller subspaces of the Hilbert space due to symmetry requirements. For example, a macroscopic atomic ensemble of N two level atoms is a collective system of N spin-$\frac{1}{2}$ systems (qubits), in a 2^N dimensional Hilbert space $\mathcal{H} = (C^2)^{\otimes N}$. However, when the dynamics of the atomic ensemble is governed by collective operations - which do not address the atoms individually - the atoms within the system are completely symmetric with respect to interchange. This class of permutation symmetric states, labeled by total spin $J = \frac{N}{2}$ (maximal value in the addition of N spin-$\frac{1}{2}$ angular momenta), is restricted [6] to a $N+1$ dimensional subspace $\mathcal{H}_{\rm Sym} = {\rm Sym}\,(C^2)^{\otimes N}$ (here 'Sym' denotes symmetrization). The eigen states $\{|J = N/2, M\rangle;\ -N/2 \leq M \leq N/2\}$ of total angular momentum operator $\hat{\vec{J}} = \frac{1}{2}\sum_{i=1}^{N} \vec{\sigma}_i$, where $\vec{\sigma}_i$ denotes Pauli spin operator of the $i^{\rm th}$ atom, span the space. A much simpler analysis of inseparability in symmetric atomic ensembles becomes possible in this space. As individual atoms are not accessible in the macroscopic ensemble, only collective measurements are feasible and any test of entanglement requiring individual control of atoms cannot be implemented experimentally. For example, spin squeezing [7], i.e., reduction of quantum fluctuations in one of the spin component orthogonal to the mean spin direction below the fundamental noise limit $N/4$ is an important *collective signature* of entanglement in symmetric N qubit systems, a consequence of two-qubit pairwise entanglement [8]. Recently [9], necessary and sufficient conditions for pairwise entanglement have been formulated in terms of negativity of intergroup covariance matrix. These are equivalent to the generalized spin squeezing inequalities [10] (for two qubit entanglement) involving collective first and second order moments of total angular momentum operator. Further, genuine three particle entanglement in symmetric multiqubit systems is shown [10] to obey inequalities involving bulk observables up to third order in total angular momentum $\hat{\vec{J}}$.

In this paper, a family of sufficient conditions to detect entanglement of atoms in a macroscopic ensemble, through collective measurements on the system, is derived. These are formulated in terms of covariance matrix condition involving averages of SO(3) irreducible tensor operators $\hat{\tau}^K_Q(N)$ of rank $K = 1, 2, \ldots N$, constructed from the total angular momentum operator $\hat{\vec{J}}$ and are identified via an arbitrary bipartite split of the symmetric states of N atoms. The physical significance of these operators are that they express $K = 1$, dipole-like; $K = 2$, quadrupole-like correlations etc., among the multiqubits. The advantages of this procedure are mainly two-fold: one, the simplicity in dealing with a large variety of correlations in multiqubit systems, and two, the entanglement conditions expressed in terms of experimentally observable signatures associated with the correlations among irreducible tensor operators. This elegant formalism enlarges the scope of the covariance matrix condition beyond those given in [9, 11]. The significance of this new approach is substantiated through illustrative examples.

The SO(3) spherical tensor operators $\hat{\tau}^K_Q(N)$ are constructed such that [12, 13] their matrix elements in the basis $\{|N/2, M\rangle\}$ are given by $\langle N/2,\, M'|\hat{\tau}^K_Q(N)|N/2,\, M\rangle = \sqrt{2K+1}\,C(N/2\,K\,N/2; MQM')$, in terms of the Clebsch Gordan (CG) coefficients [14]. These irreducible tensors are orthogonal, ${\rm Tr}(\hat{\tau}^K_Q(N)\,\hat{\tau}^{K'}_{Q'}(N)^\dagger) = (N+1)\,\delta_{K,\,K'}\delta_{Q,\,Q'}$, and the set $\{\hat{\tau}^K_Q(N);\ K = 0, 1, 2, \ldots N,\ -K \leq Q \leq K\}$ forms a linearly independent basis of operators in the Hilbert space of spin $J = N/2$ states. Thus a useful representation for the density operator of symmetric N-qubits is given in terms of these operators by

$$\hat{\rho}(N) = \frac{1}{(N+1)} \sum_{K=0}^{N} \sum_{Q=-K}^{K} \hat{\tau}_Q^{K\dagger}(N)\; t_Q^K(N). \tag{1}$$

It is completely specified by $(N+1)^2 - 1$ irreducible tensor moments,

$$t_Q^K(N) = \mathrm{Tr}[\hat{\rho}_{\mathrm{sym}}(N)\, \hat{\tau}_Q^K(N)] = (-1)^Q\, t_{-Q}^{K*}(N), \tag{2}$$

with $t_0^0(N) = 1$ because $\mathrm{Tr}\hat{\rho} = 1$. An important *composition law* appropriate for examining multipole correlations between two constituent symmetric parts of this N particle system, characterized by angular momenta $j_1 = N_1/2$, and $j_2 = N_2/2$ (with $N = N_1 + N_2$), of the ensemble is constructed in terms of direct product of spherical tensors $\tau_q^{\kappa}(N_1) \otimes \tau_{q'}^{\kappa'}(N_2)$:

$$\hat{\rho}(N_1, N_2) = \frac{1}{(N_1+1)(N_2+1)} \sum (\hat{\tau}_q^{\kappa\dagger}(N_1) \otimes \hat{\tau}_{q'}^{\kappa'\dagger}(N_2))\, t_{qq'}^{\kappa\kappa'}(N_1, N_2), \tag{3}$$

where $t_{qq'}^{\kappa\kappa'}(N_1, N_2) = \mathrm{Tr}[\hat{\rho}(N_1, N_2)\, \hat{\tau}_q^{\kappa}(N_1) \otimes \hat{\tau}_{q'}^{\kappa'}(N_2)] = (-1)^{q+q'}\, t_{-q\,-q'}^{\kappa\kappa'\,*}(N_1, N_2)$; $\kappa = 0, 1, \ldots, N_1$, $\kappa' = 0, 1, \ldots, N_2$; $-\kappa \leq q \leq \kappa, -\kappa' \leq q' \leq \kappa$. Equations (1) and (3) represent the N-qubit symmetric system in two equivalent ways and thus the tensor parameters appearing therein are related as will be shown presently. These relations form the central core of the theory presented here.

By taking trace over N_2 particles from the composite system (using the representation (3)), we obtain the density matrix of N_1 particles. We find that the tensor parameters $t_q^{\kappa}(N_1)$ characterizing the N_1 subsystem are given by,

$$t_q^{\kappa}(N_1) = \mathrm{Tr}\left(\hat{\rho}(N_1)\, \hat{\tau}_q^{\kappa}(N_1)\right) = t_{q0}^{\kappa 0}(N_1, N_2); \quad \kappa = 0, 1, \ldots, N_1. \tag{4}$$

Similar consideration for N_2-subsystem leads to $t_{q'}^{\kappa'}(N_2) = t_{0q'}^{0\kappa'}(N_1, N_2)$. The second set of relations connecting the tensor parameters of the bipartite system given by (3) with those of (1), are obtained by using the orthogonality property of the tensor operators:

$$t_{qq'}^{\kappa\kappa'}(N_1, N_2) = \sum_{K,Q} \mathcal{F}_{qq';\,KQ}^{(\kappa\kappa')}(N_1, N_2, N)\, t_Q^K(N), \tag{5}$$

where $|\kappa - \kappa'| \leq K \leq \kappa + \kappa'$; $-K \leq Q \leq K$, and $\mathcal{F}_{qq';\,KQ}^{(\kappa\kappa')}(N_1, N_2, N)$ are found to be

$$\begin{aligned} \mathcal{F}_{qq';\,KQ}^{(\kappa\kappa')}(N_1, N_2, N) &= \mathrm{Tr}\left((\hat{\tau}_q^{\kappa}(N_1) \otimes \hat{\tau}_{q'}^{\kappa'}(N_2)\, \hat{\tau}_Q^K(N)\right) \\ = \left[\tfrac{N_1}{2}\right] \left[\tfrac{N_2}{2}\right] \left[\tfrac{N}{2}\right] [\kappa]\, [\kappa']\, &C(\kappa\, \kappa'\, K; q\, q'\, Q) \left\{ \begin{array}{ccc} \frac{N_1}{2} & \frac{N_2}{2} & \frac{N}{2} \\ \frac{N_1}{2} & \frac{N_2}{2} & \frac{N}{2} \\ \kappa & \kappa' & K \end{array} \right\}. \end{aligned} \tag{6}$$

Here '{ }' denotes the Wigner-9j symbol [12, 14] and $[a] = \sqrt{2a+1}$. Using the properties of the 9j-symbol and the CG coefficients [12] for the special values $\kappa = 0, q = 0$ in (5) and (6), we obtain

$$t_{q0}^{\kappa 0}(N_1, N_2) = \mathcal{P}_{\kappa}(N_1, N_2)\, t_q^{\kappa}(N), \tag{7}$$

where $\mathcal{P}_{\kappa}(N_1, N_2) = \frac{N_1!}{(N)!} \sqrt{\frac{(N_1+1)(N+\kappa+1)!(N-\kappa)!}{(N+1)(N_1+\kappa+1)!(N_1-\kappa)!}}$; $\kappa = 0, 1, 2, \ldots N_1$. By replacing $N_2 \to N_2 - N_1$, (where $N_2 \geq N_1$) in both sides of (7) and using (4), we obtain an important equivalent relation

$$t_q^{\kappa}(N_1) = \mathcal{P}_{\kappa}(N_1, N_2 - N_1)\, t_q^{\kappa}(N_2). \tag{8}$$

The product tensor parameters $t_{qq'}^{\kappa\kappa'}(N_1, N_2)$ of the bipartite system exhibit a similar relationship with the corresponding coefficients $t_{qq'}^{\kappa\kappa'}(\kappa, \kappa')$ of the $\kappa + \kappa'$ subsystem. This follows by expressing $t_Q^K(N)$ of the composite system in the RHS of (5) in terms of $\kappa + \kappa'$ subsystem parameters $t_Q^K(\kappa + \kappa')$ (obtained by substituting $N_1 = \kappa + \kappa', N_2 = N - (\kappa + \kappa')$ in (8)). Alternately, we can also relate $t_{qq'}^{\kappa\kappa'}(N_1, N_2)$ to $t_Q^K(\kappa + \kappa')$ by choosing $N_1 = \kappa$, $N_2 = \kappa'$ in (5). Comparing the resulting equations for $t_{qq'}^{\kappa\kappa'}(N_1, N_2)$ and $t_{qq'}^{\kappa\kappa'}(\kappa, \kappa')$ and using explicit expressions for the associated 9j-symbols [12] we obtain, after some algebraic manipulation,

$$\begin{aligned} t_{qq'}^{\kappa\kappa'}(N_1, N_2) &= f(N_1, \kappa)\, f(N_2, \kappa')\, t_{qq'}^{\kappa\kappa'}(\kappa, \kappa'), \\ f(N_\alpha, \kappa) &= \sqrt{\frac{(N_\alpha+1)(2\kappa+1)!}{(\kappa+1)(N_\alpha+\kappa+1)!(N_\alpha-\kappa)!}}; \quad \alpha = 1, 2. \end{aligned} \tag{9}$$

Equations (5), (8) and (9) prove to be significant in identifying collective signatures of entanglement in symmetric atomic ensembles, obtained through an analysis of the bipartite representation (3).

Consider a set of $2\,(2\kappa + 1)$ operators $\hat{A}_q^{\kappa} = \hat{\tau}_q^{\kappa}(N_1) \otimes \hat{I}_{N_2}$ and $\hat{B}_q^{\kappa} = \hat{I}_{N_1} \otimes \hat{\tau}_q^{\kappa}(N_2)$ (where $\hat{I}_{N_i} = \hat{\tau}_0^0(N_i)$ denotes the identity operator). Arranging them as a column $\xi^{(\kappa)}$ (corresponding row of operators being $\xi^{(\kappa)} = (\hat{A}_q^{\kappa\dagger}, \hat{B}_q^{\kappa\dagger})$, define the 2κth order covariance matrix for the symmetric system as

$$V_{ij}^{(2\kappa)} = \frac{1}{2} \left\{ \Delta\xi_i^{(\kappa)},\, \Delta\xi_j^{(\kappa)} \right\}, \tag{10}$$

where $\Delta\xi^{(\kappa)} = \xi^{(\kappa)} - \langle \xi^{(\kappa)} \rangle$ and $\{\Delta\xi_i^{(\kappa)},\, \Delta\xi_j^{(\kappa)}\} = \Delta\xi_i^{(\kappa)}\, \Delta\xi_j^{(\kappa)} + \Delta\xi_j^{(\kappa)}\, \Delta\xi_i^{(\kappa)}$. $V^{(2\kappa)}$ exhibits a $(2\kappa + 1) \times (2\kappa + 1)$ block matrix form, $V^{(2\kappa)} = \begin{pmatrix} A^{(2\kappa)}(N_1) & C^{(2\kappa)}(N_1, N_2) \\ C^{(2\kappa)\dagger}(N_1, N_2) & B^{(2\kappa)}(N_2) \end{pmatrix}$. The diagonal blocks $A_{qq'}^{(2\kappa)}(N_1) = \frac{1}{2} \langle \{\Delta\hat{A}_q^{\kappa}, \Delta\hat{A}_{q'}^{\kappa}\} \rangle$, $B_{qq'}^{(2\kappa)}(N_2) = \frac{1}{2} \langle \{\Delta\hat{B}_q^{\kappa}, \Delta\hat{B}_{q'}^{\kappa}\} \rangle$ correspond to multipole correlations among the intra-group tensors and the off-diagonal block $C_{qq'}^{(2\kappa)}(N_1, N_2) = \frac{1}{2} \langle \{\Delta\hat{A}_q^{\kappa}, \Delta\hat{B}_{q'}^{\kappa}\} \rangle$ comprises of inter-group multipole correlations. Explicitly,

$$C_{qq'}^{(2\kappa)}(N_1, N_2) = (-1)^{q'}\, [t_{q\,-q'}^{\kappa\kappa}(N_1, N_2) - t_{q0}^{\kappa 0}(N_1, N_2)\, t_{0\,-q'}^{0\kappa}(N_1, N_2)]. \tag{11}$$

Here we focus on the $(2\kappa + 1) \times (2\kappa + 1)$ hermitian cross-correlation matrix $C^{(2\kappa)}$ and prove the following theorem:

Theorem : The cross-correlation matrix $C^{(2\kappa)}(N_1, N_2)$ of a given rank κ associated with any partition (N_1, N_2) of an N-qubit symmetric system is necessarily positive semidefinite for all separable symmetric N-qubit bipartite states. The sign of the corresponding matrix $C^{(2\kappa)}(\kappa, \kappa)$, associated with the 2κ atom reduced system with equal partition, suffices to determine that of $C^{(2\kappa)}(N_1, N_2)$, irrespective of the partitioning.

Proof: In the product representation (3) with an arbitrary partition (N_1, N_2), a separable symmetric N-qubit state has the following structure:

$$\hat{\rho}_{\mathrm{sep}} = \sum_w p_w\, \hat{\rho}_w(N_1) \otimes \hat{\rho}_w(N_2), \quad 0 \leq p_w \leq 1; \quad \sum_w p_w = 1 \tag{12}$$

where $\hat{\rho}_w(N_i)$ denotes the density matrices of the subensemble of N_i qubits and is expressible as (1), in terms of $\hat{\tau}^{\kappa}_q(\hat{N}_i)$: $\hat{\rho}_w(N_i) = \frac{1}{(N_i+1)} \sum_{\kappa=0}^{N_i} \sum_{q=-\kappa}^{\kappa} \hat{\tau}^{\kappa\dagger}_q(N_i)\, t^{\kappa}_q(N_i, w)$. In a separable symmetric state (12) we have,

$$\begin{aligned} t^{\kappa\kappa}_{qq'}(N_1, N_2) &= \mathrm{Tr}\left(\hat{\rho}_{\rm sep}\, [(\hat{\tau}^{\kappa}_q(N_1) \otimes \hat{\tau}^{\kappa}_{q'}(N_2)]\right) \\ &= \sum_w p_w\, \mathrm{Tr}\left(\hat{\rho}_w(N_1)\, \hat{\tau}^{\kappa}_q(N_1)\right) \mathrm{Tr}\left(\hat{\rho}_w(N_2)\, \hat{\tau}^{\kappa}_{q'}(N_2)\right) \\ &= \sum_w p_w\, t^{\kappa}_q(N_1, w)\, t^{\kappa}_{q'}(N_2, w). \end{aligned} \tag{13}$$

Without any loss of generality, we may assume that $N_2 \geq N_1$. Now, employing (7) we express $t^{\kappa}_{q'}(N_2, w) = [\mathcal{P}_\kappa(N_1, N_2 - N_1)]^{-1}\, t^{\kappa}_q(N_1, w)$, which leads to an interesting form for the matrix $C^{(2\kappa)}(N_1, N_2)$ of (11) in a separable symmetric state:

$$C^{(2\kappa)}_{q\,q'}(N_1, N_2) = [\mathcal{P}_\kappa(N_1, N_2 - N_1)]^{-1}\, C^{(2\kappa)}(N_1, N_1) \tag{14}$$

Consider an hermitian quadratic form

$$Q^{\kappa} = X^{\kappa\dagger}\, C^{(2\kappa)}(N_1, N_2) X^{\kappa} = \sum_{q,\ q'=-\kappa}^{\kappa} C^{(2\kappa)}_{qq'}(N_1, N_2)\, X^{\kappa *}_q\, X^{\kappa}_{q'}, \tag{15}$$

with $X^{\kappa} \in R^{(2\kappa+1)}$ being an arbitrary real column vector, whose spherical components [12] are denoted by $X^{\kappa}_q = (-1)^q\, X^{\kappa *}_{-q}$. In a separable symmetric state (12) it is readily seen that $Q^{\kappa}_{\rm sep} = [\mathcal{P}_\kappa(N_1, N_2 - N_1)]^{-1}\, \Pi^{\kappa}_{\rm sep} \geq 0$, with $\Pi^{\kappa}_{\rm sep} = X^{\kappa\dagger}\, C^{(2\kappa)}(N_1, N_1) X^{\kappa} = \sum_w p_w [\sum_q X^{\kappa *}_q\, t^{\kappa}_q(N_1, w)]^2 - [\sum_w p_w \sum_q X^{\kappa *}_q\, t^{\kappa}_q(N_1, w)]^2$, a positive semidefinite quantity. This proves the first part of our theorem.

The second part of the theorem follows from (9), leading to the result

$$C^{(2\kappa)}(N_1, N_2) = f(N_1, \kappa)\, f(N_2, \kappa)\, C^{(2\kappa)}(\kappa, \kappa), \tag{16}$$

i.e., the covariance matrix $C^{(2\kappa)}(N_1, N_2)$ of an arbitrary symmetric system is proportional (with an overall positive multiplication factor) to that associated with the equal partitioning of a 2κ qubit reduced system. (Note that the covariance matrix $C^{(2\kappa)}(N_1, N_2)$ could be related to its reduced system counterpart $C^{(2\kappa)}(n_1, n_2)$, which is then seen to be proportional to $C^{(2\kappa)}(n_1, n_1)$; $n_1 \leq n_2$. Further, $C^{(2\kappa)}(n_1, n_1)$ may be related to $C^{(2\kappa)}(n, n)$; $n < n_1$ etc. This can go down all the way upto $C^{(2\kappa)}(\kappa, \kappa)$. Hence, the positivity (negativity) of $C^{(2\kappa)}(N_1, N_2)$ has its origin in the 2κ-qubit covariance matrix $C^{(2\kappa)}(\kappa, \kappa)$, with equal partition (κ, κ).) □

Thus, for any arbitrary (pure or mixed) symmetric ensemble of qubits, $C^{(2\kappa)}(\kappa, \kappa) < 0$, for various orders $\kappa = 1, 2, \ldots; 2\kappa \leq N$, serves as a sufficient condition of entanglement and leads to a family of inseparability conditions associated with the quantum correlations between inter-group tensor operators i.e., $\langle \Delta \hat{A}^{(\kappa)}_q\, \Delta \hat{B}^{(\kappa)}_q \rangle$. As the product tensor parameters $t^{\kappa\kappa}_{qq'}(\kappa, \kappa)$ and $t^{\kappa 0}_{q0}(\kappa, \kappa)$, $t^{0\kappa}_{0q})(\kappa, \kappa)$ - specifying the covariance matrix $C^{(2\kappa)}(\kappa, \kappa)$ - are given in terms of collective tensor moments $t^{K}_{Q}(N)$, (see (5), (8) and (9)), the negativity of the covariance matrix $C^{(2\kappa)}(\kappa, \kappa)$ is readily expressed in terms of averages of symmetrized homogeneous $2\kappa^{\rm th}$ order polynomials [13] of $\hat{\vec{J}}$, thus leading to a family of *multipole-like collective signatures* of entanglement.

We now show that the spin squeezing inequality [9, 10] is a consequence of the intergroup dipole correlations (for $\kappa = 1$) viz., $C^{(2)}(N_1, N_2) < 0$: In the 3×3 matrix $C^{(2)}(N_1, N_2)$ we substitute $t^{\kappa\kappa'}_{qq'}(N_1, N_2); \kappa, \kappa' = 0, 1$ in terms of total system collective parameters $t^{K}_{Q}(N)$ (see (5)) using explicit values [12] for Wigner $9j$ symbols and CG coefficients. Following this by a unitary transformation corresponding to a change from spherical basis [12] $\mathbf{e}_\mu,\ \mu = \pm 1, 0$ to Cartesian basis $\mathbf{e}_i,\ i = x, y, z$ leads to

$$U\, C^{(2)}(N_1, N_2)\, U^{\dagger} = \mathcal{A}\, [-\frac{N}{4}\, \mathcal{I} + V + \frac{1}{N}\, S S^{T}] \tag{17}$$

(where $\mathcal{I}$ denotes the 3×3 identity matrix; $V_{\alpha\,\beta} = \frac{1}{2} \left\langle (\hat{J}_\alpha \hat{J}_\beta + \hat{J}_\beta \hat{J}_\alpha) \right\rangle - \langle \hat{J}_\alpha \rangle \langle \hat{J}_\beta \rangle$; $S_\alpha = \langle \hat{J}_\alpha \rangle$ and $\mathcal{A} = \frac{12}{N(N-1)} \sqrt{\frac{N_1\, N_2}{(N_1+2)(N_2+2)}}$). Hence, $C^{(2)}(N_1, N_2) < 0 \Longleftrightarrow U\, C^{(2)}(N_1, N_2)\, U^{\dagger} < 0 \Longleftrightarrow V + \frac{1}{N}\, S S^{T} < \frac{N}{4}$, which is the known spin squeezing inequality [9, 10], deduced here from any bipartite division (N_1, N_2). (The covariance matrix condition of [11] coincides with that given here *only* for dipole correlations.)

An important application of our theorem concerns the Dicke states [15] $|\frac{N}{2}, M\rangle$; $-\frac{N}{2} \leq M \leq \frac{N}{2}$. These states provide an excellent set of physically relevant multiatom symmetric states for illustrating other types of multipole correlations. The collective tensor moments for the Dicke states are $t^{K}_{Q}(N) = \langle \frac{N}{2}, M | \hat{\tau}^{K}_{Q}(N/2) | \frac{N}{2}, M \rangle = [\kappa]\, C(\frac{N}{2}\, K\, \frac{N}{2}; M\, 0\, M)\, \delta_{Q\,0}$. This corresponds to non-zero product tensor parameters $t^{\kappa\kappa'}_{q-q}(N_1, N_2)$ (obtained from (5)) implying that $C^{(2\kappa)}(N_1, N_2)$ matrix is diagonal (see the definition (11)). Except for $|\frac{N}{2}, \pm\frac{N}{2}\rangle$, which are product states, the covariance matrix $C^{(2\kappa)}(\kappa, \kappa)$ of each of the Dicke states is negative, for all orders κ, showing the quantum nature of multipole-like correlations of various orders. Recently [16], an experimental scheme to reconstruct the spin-excitation number distribution of the collective spin states (i.e., tomographic reconstruction of the diagonal elements of the density matrix in the Dicke basis) of macroscopic ensembles containing $\sim 10^{11}$ atoms, with low mean spin excitations, has been proposed. Implementation of such schemes would enable experimental detection of *collective quantum multipole-like correlations* in macroscopic assembly of entangled atoms.

As an illustration of our method, we consider symmetric mixed states of the form

$$\hat{\rho}_i = \frac{(1-x)}{(N+1)}\, \mathcal{I} + x\, |\phi_i\rangle\, \langle\phi_i|; \quad 0 \leq x \leq 1, \tag{18}$$

where $\mathcal{I}$ denotes $(N+1) \times (N+1)$ unit matrix; the states $|\phi_i\rangle$, for $i = 1, 2, 3$, are given by $|\phi_1\rangle = |\frac{N}{2}, \frac{N}{2} - 1\rangle$, $|\phi_2\rangle = |\frac{N}{2}, 0\rangle$ and $|\phi_3\rangle = \frac{1}{\sqrt{2}}(|\frac{N}{2}, \frac{N}{2}\rangle + |\frac{N}{2}, -\frac{N}{2}\rangle)$. We find the values of x, - using the condition $C^{(2\kappa)}(N_1, N_2) < 0$ - as a function of number of atoms, for which $\hat{\rho}_i$ of (18) are inseparable. For the state $\hat{\rho}_3$ we find that $C^{(2\kappa)}(\kappa, \kappa)$, are all positive for $2\kappa < N$ and the highest order co-variance matrix $C^{(N)}(N/2, N/2) < 0$.



FIG. 1: Threshold value $x_{\min}$, evaluated from the multipole inseparability condition $C^{(2\kappa)}(\kappa, \kappa) < 0$, with $\kappa = 1$ to $\kappa = 5$ (uppermost curve to the lower ones respectively) for the states $\hat{\rho}_1$ (see (a)) and $\hat{\rho}_2$ (see (b)), as a function of the number of atoms N. (c): $x_{\min}$ obtained from the negativity of the highest order covariance matrix $C^{(N)}(N/2, N/2)$ for the states $\hat{\rho}_1$ ($'+'$), $\hat{\rho}_2$ ($'\Box'$) and $\hat{\rho}_3$ ($'\times'$).

These results are presented in graphical form in Fig. 1. From Fig. 1(a) and 1(b) it is clear that dipole ($\kappa = 1$) quantum correlations lead to $x_{\min} \to 1$ for large N values, implying that the mixed states $\hat{\rho}_i;\ i = 1, 2$ are separable throughout the range $0 \leq x < 1$ in this limit. However, higher order multipole correlations are more effective in revealing that these states are indeed entangled over a larger domain of x. The range of inseparability is sensitive to the difference $N_e \sim N_g$ of the number of atoms in ground and excited states (which is $N - 1$ in $|\phi_1\rangle$ and *zero* in $|\phi_2\rangle$). Further, from Fig. 1(c) we find that highest order quantum correlations lead to $x_{\min} \to 0$ in the large N limit, implying that all the three mixed states $\hat{\rho}_i$, for $i = 1, 2, 3$, are entangled in the range $0 < x \leq 1$, when $N \to \infty$.

In conclusion, we have shown here that SO(3) irreducible tensor representation provides a powerful method to investigate inseparability in symmetric multiqubit systems. A family of sufficient conditions of inseparability to detect multipole-like *collective quantum correlations* derived here may be useful for experimental characterization of entanglement in macroscopic atomic ensembles. These conditions are generalizations of the ones obtained in [9, 11]. More specifically, instead of the Cartesian tensor product observables of Ref. [11], SO(3) irreducible tensor observables are shown here to characterize any bipartite division (N_1, N_2) of symmetric N-qubit systems. The two techniques serve different purposes. The first considers groups of qubits to investigate the intergroup entanglement, whereas the second describes them in terms of spherical tensors involving collective N-qubit angular momenta. The latter sheds light on entanglement among dipole-like, etc. multiqubit correlations, which may be physically observable [16] in macroscopic ensembles of symmetric atoms. An important consequence is that the Dicke states exhibit quantum multipole-like correlations of all orders. Moreover, our approach is directly applicable to characterize entanglement in spatially separated bipartite symmetric atomic ensembles, for example, two macroscopic gas samples of cesium atoms [17].

---

[*] Electronic address: arutth@rediffmail.com

[1] J. Hald, J. L. Sørensen, C. Schori, and E. S. Polzik, Phys. Rev. Lett. **83**, 1319 (1999); A. Kuzmich, L. Mandel, and N. P. Bigelow, Phys. Rev. Lett. **85**, 1594 (2000)

[2] D. J. Wineland, J. J. Bolinger, W. M. Itano, and D. J. Heinzen, Phys. Rev. A **50**, 67 (1994).

[3] B. Yurke, Phys. Rev. Lett. **56**, 1515 (1994); M. Kitagawa, and M. Ueda, Phys. Rev. Lett. **56**, 1852 (1991); V. Giovannetti, S. Lloyd, and L. Maccone, Science **306**, 1330 (2004); C. Lee, Phys. Rev. Lett. **97**, 150402 (2006).

[4] H. J. Briegel, W. Dur, J. I. Cirac, P. Zoller, Phys. Rev. Lett. **81**, 5932 (1998); B. Julsgaard, J. Sherson, J. I. Cirac, J. Fiurasek, and E. S. Polzik, Nature **432**, 482 (2004); J. F. Sherson, H. Krauter, R. K. Olsson, B. Julsgaard, K. Hammerer, J. I. Cirac and E. S. Polzik, Nature **443**, 557 (2006).

[5] K. G. H. Vollbrecht, and R. F. Werner, Phys. Rev. A **64**, 062307 (2001); H. P. Breuer, Phys. Rev. A **71**, 062330 (2005); J. Phys. A **38**, 9019 (2005); D. Chruściński and A. Kossakowski, Phys. Rev. A **73**, 062314, 062315 (2006).

[6] J. K. Stockton, J. M. Geremia, A. C. Doherty, and H. Mabuchi, Phys. Rev. A **67**, 022112 (2003); J. M. Geremia, J. K. Stockton, and H. Mabuchi, Phys. Rev. A **73**, 042112 (2006).

[7] M. Kitagawa, and M. Ueda, Phys. Rev. A **47**, 5138 (1993).

[8] A. Sørensen, L.-M. Duan, J. I. Cirac, and P. Zoller, Nature (London) **409**, 63 (2001); X. Wang, and B. C. Sanders, Phys. Rev. A **68**, 012101 (2003); A. R. Usha Devi, M. S. Uma, R. Prabhu, and Sudha, Int. J. Mod. Phys. B **20**, 1917 (2006).

[9] A. R. Usha Devi, M. S. Uma, R. Prabhu, and A. K. Rajagopal, Phys. Lett. A **364**, 203 (2007).

[10] J. K. Korbicz, J. I. Cirac, and M. Lewenstein, Phys. Rev. Lett. **95**, 120502 (2005); **95**, 259901(E) (2005); J. K. Korbicz *et. al.*, Phys. Rev. A **74**, 052319 (2006).

[11] A. R. Usha Devi, R. Prabhu, and A. K. Rajagopal, Phys. Rev. Lett. **98**, 060501 (2007).

[12] D. A. Varshalovich, A. N. Moskalev, and V. K. Khersonskii, *Quantum Theory of Angular Momentum* (World Scientific, Singapore, 1988).

[13] The spherical tensors $\hat{\tau}^K_Q$ are homogeneous polynomials of order K constructed from the angular momentum operators $\hat{\vec{J}}$ as, $\hat{\tau}^K_Q = \mathcal{N}_{KJ}\,(\hat{\vec{J}} \cdot \vec{\nabla})^K\ \mathcal{Y}_{KQ}(x, y, z)$ where $\mathcal{Y}_{KQ}(x, y, z) = r^K\, Y_{KQ}(\theta, \phi)$ are the solid harmonic functions. The choice of normalization factor $\mathcal{N}_{KJ} = \frac{2^K}{K!}\sqrt{\frac{4\pi\,(2J-K)!}{(2J+1)(2J+K+1)!}}$ leads to their matrix elements in the $\{|J\,M\rangle\}$ basis as given in the text. Note that the operators $\hat{\tau}^K_Q$ are related to the $\hat{T}_{KQ}$ of Ref. [12] through $\hat{\tau}^K_Q = [J]\,\hat{T}_{KQ}$, so that $\hat{\tau}^0_0$ is a $(2J+1) \times (2J+1)$ identity matrix.

[14] We mainly follow the notations and conventions of M. E. Rose, *Elementary Theory of Angular Momentum* (John

Wiley, New York, 1957).
[15] R. H. Dicke, Phys. Rev. **93**, 99 (1954).
[16] K. Usami, J. Takahashi, and M. Kozuma, Phys. Rev. A **74**, 043815 (2006).
[17] B. Julsgaard, A. Kozhekin, and E. S. Polzik, Nature **413**, 400 (2001).